Relativistic nanophotonics: creating extreme plasma conditions and fields with ultrafast laser pulses

Jorge J. Rocca

Colorado State University, Fort Collins, Colorado 80523, USA jorge.rocca@colostate.edu

R. Hollinger¹, V. N Shlyaptsev¹, M.G. Capeluto¹, A, Curtis¹, A. Moreau¹, C, Calvi¹, S. Kardorf¹, V. Kaymak², A. Pukhov², Y. Wang¹, S, Wang¹, A. Rockwood¹, C.S. Menoni¹

¹ CSU, Fort Collins, Colorado, ² Heinrich-Heine-Universität Düsseldorf






LaserNetUS






High Power Laser and ultra-Intense Laser-Matter Interaction at CSU's Advanced Beam Laboratory






Path to extreme plasma conditions by irradiation of aligned nanowire arrays






Sun Core 240 Gbar

NIF Implosion >150 Gbar




Nanowire array plasma $I = 1 \times 10^{22} \text{ W cm}^{-2}$ > 300 Gbar (predicted)




Volumetric heating of UHED plasmas by Irradiation of aligned nanowire arrays with utrashort laser pulses of relativistic intensity





Laser irradiation of aligned nanowire arrays at I =5x 10 ¹⁸ W cm-² creates multi-KeV temperature plasmas with densities up to 100X critical






M. Purvis et al. *Nature Photonics*, **7**,796, (2013)

Nanowire arrays SEM Images






At $5 \times 10^{18} \text{ W cm}^{-2}$ Nickel He- α emission exceeds K- α




0.5 J pulse , 55 fs duration



M. Purvis et al. Nature Photonics, 7, 796 (2013)

Measurement of heat penetration depth monitoring emission from buried tracer element shows volumetric heating






Absorption of intense ultrashort laser pulses in nanowire arrays








Irradiation with two delayed ultrashort pulse determines the time available for efficient energy coupling






- 80nm Ni wires 12% solid density
- Pulse intensities 4x10¹⁹ -1x10²⁰ Wcm⁻²

Energy density and pressure distribution

55 nm diameter, Ni wires, 13% solid density; $I = 4x10^{19} \text{ W cm}^{-2}$, 55 fs






C. Bargsten et al. Science Advances, 3, e1601558 (2017),

Matter in the ultra high-energy density regime




(> 1x10⁸ J cm⁻³, pressures > 1 Gbar)



M. Purvis et al. *Nature Photonics*, **7**, 796 (2013)

In volumetrically heated nanoarray plasmas radiation cooling can dominate over hydrodynamic cooling






Increased X-ray CE from nanowires (< 1 % efficiency): G Kultzar et al. (PRL, 2000); T. Nishikawa et al (Appl. Phys.B, 2004); S. Modal et al (Phys. Rev.B, 2011)

Record Conversion Efficiency into > 1 KeV X-rays



CE ~ 20 percent, with individual shots up to 24% CE into 4π





R. Hollinger et al., Optica, **4**, 1344, (2017)


Single Shot X-ray Imaging (3 µm Spatial Resolution)



17








APS DPP 2015

Deuterium ion acceleration and neutron production in deuterated nanowires






Deuteron energies up to 3 MeV measured

Colorado State University

Intensity 8 x 10¹⁹ W cm⁻²




Nanowire




Deuteron and Neutron Detection Setup





Neutron Time of Flight: D-D Fusion Neutrons






A. Curtis et al, Nature Comm., 1077 (2018)

Nanowire arrays display 500 X increased neutron yield as compared with CD₂ flat targets






Maximum measured neutrons/shot: (>2.5 x 10 6 neutrons/Joule)


Predicted Formation of Relativistic Nano-pinch





Nanowire pinch

Laser-induced return current densities of ~ 0.1 Giga-Amperes per μm^2 through the wires




V. Kaymak et al, Phys. Rev. Lett,.117, 035004 (2016)

The large return current generates a Giga-Gauss azimuthal magnetic field and nano-scale Z-pinch with > 1000 times ne_c



Carbon nanowires 300 nm diameter, I=5 x10²¹ Wcm⁻²




V. Kaymak et al, Phys. Rev. Letters 117, 035004 (2016).

Self-generated quasi-static azimuthal magnetic field pinches nanowires into hot plasmas with $n_e > 9x10^{24}$ cm⁻³, > 1000 times the critical density




Electron density maps 300 nm C nanowires, I= 5x10²¹W cm⁻²



V. Kaymak et al, Phys. Rev. Letters 117, 035004 (2016).

Au atoms ionized 69 times in an electron beam ion trap (EBIT)





Densities > 10 orders of magnitude below solid density


G.V. Brown et al Physical Review E 77, 066406 (2008)

High energy lasers lonizes gold to Au⁺⁵⁷



OMEGA LASER (U. Rochester)







 E_L = 9kJ ; T_e =6.5 keV

Previous experiments at 5x10¹⁸ W cm⁻³ show emission from up to Ni-like Au⁵²⁺ and optical volumetric heating






M. Purvis et al Nature Photonics 7, 796- (2013)



Irradiation intensity of 1 x 10 ²² W cm⁻³ predicted to create extreme ionization states in Au (eg. F-like Au)



Ion charge distribution as a function of depth:



3.3 Hz, 0.85 PW- Ti:Sapphire Laser System





3828 Vol. 42, No. 19 / October 1 2017 / Optics Letters

Letter

Optics Letters

0.85 PW laser operation at 3.3 Hz and high-contrast ultrahigh-intensity λ = 400 nm second-harmonic beamline

YONG WANG, 1.1.* SHOUJUN WANG, 1.† ALEX ROCKWOOD, 2.† BRADLEY M. LUTHER, 3 REED HOLLINGER, 1 ALDEN CURTIS, 1 CHASE CALVI, 2 CARMEN S. MENONI, 1.3 AND JORGE J. ROCCA 1.2

¹Electrical and Computer Engineering Department, Colorado State University, Fort Collins, Colorado 80523, USA

²Physics Department, Colorado State University, Fort Collins, Colorado 80523, USA


³Chemistry Department, Colorado State University, Fort Collins, Colorado 80523, USA

*Corresponding author: yongw@colostate.edu



ALEPH: Advanced Laser for Extreme Photonics








"I saw a small iridescent sphere of almost unbearable brilliance."

JLB


Slab amplifier on double-pass configuration 18 J at 3.3 Hz






Pump laser slab amplifiers for 36 J Ti:Sapphire amplifier operating at 3.3 Hz repetition rate











3.3 Hz laser run, 36.3 J average energy before compressor






- ☐ Average pulse energy 36.3 J
- ☐ Shot-to-shot variation ~1.7% rms



Spatial beam profile

Laser spectra, 30 fs pulse duration







Autocorrelation trace of the compressed pulses obtained from a single shot real-time spectral phase measurement.


Ultra-high contrast pulses: > 1x10¹²

Frequency Doubling and filtering of fundamental beam









f/2 parabola achieves 6.5×10²¹ W/cm²






Relativistic Induced Transparency and Electron Acceleration in Dense CD₂ Nanowire array




Irradiation intensity I = 2.7x10²¹ W cm-²



Relativistically induced transparency allows for deep laser pulse propagation into overdense CD₂ plasma






Relativistically Transparent Nanowire array target generates MeV electrons with increased energy and total flux




CD₂ Nanowire Array Electron Spectra (forward direction)



- Integrated flux enhanced 5x for typical NW shot over solid
- Best shot flux enhanced 24.5x over solid target


Increase of 3.6 X in > 1 MeV Gamma ray flux





Ion Trajectories CD₂ nanowires, I= 5x10²¹ W cm⁻²






Neutron production increased to $> 2x10^7$ n/shot




TOF detector at 2m with 10 cm lead shielding



Ion trajectories




 CD_2 Nanowires, 400 nm diameter, 5 x 10 21 W cm $^{-3}$



Ion beam in laser backward directions









Angular Ion Distribution Measurement (> 13.3MeV D)





Au Nanowires






I= 4x10²¹ W cm⁻³ 15 % solid density

Laser drives hot electrons at tip of wire into plasma






- Laser drives hot electrons at tip of wire into plasma
- Peak electron
 energy density of
 >1TJ/cc at tip of
 wire
- Electrons deposit energy through the volume ending with an energy density of ~100 GJ/cc
- Large current density causes
 J x B force to pinch the wire

Au L shell spectra from solid and near solid density plasmas show emission from ions up to N-like Au⁷²⁺








- Both targets show emission up to N-like Au⁷²⁺
- Nanowire target charge state peaks around Allike Au⁶⁶⁺
- Solid target is peaks around Nelike Au⁶⁹⁺

In steady state ionization equilibrium Neon-like ions appear at temperatures > 10 KeV








Simulation of solid target heat penetration depth agrees with experiment



+500fs after peak of laser pulse









Buried Ni tracer reveals heat penetration in nanowire array exceeds 8 micrometers



 $I=4x10^{21} Wcm^{-2} (a_o = 21)$




~8µm Au on Ni 100nm 15% NW




He-Like Au⁺⁷⁷ solid density plasma predicted




 $4x10^{21}$ Wcm⁻² with a 5 μ m FWHM diameter focal spot



Scaling to $1x10^{22}$ W cm⁻³ ($a_0 = 34$)







Contributors:





Conclusions:



Interaction of highly relativistic ultrashort laser pulses with nanostructures

- Interactions of high contrast, relativistic laser pulses with nanostructures creates ultra high energy density plasmas with extreme conditions.
- Demonstrated Au plasmas with extraordinary degree of ionization, N-like Au line emission.
- Measured record conversion efficiency optical laser light into picosecond x-ray pulses of up to 20% for hv >1 keV
- Measured increased flux and energy of fast electron and ions
- Generated flashes of quasi monoenergetic neutrons from D-D fusion

Work supported by US DOE, AFOSR, ONR

